



**Huron County General Health District**  
 180 Milan Ave., Norwalk OH 44857  
 419-668-1652 Toll free 1888-694-2443  
 Fax: 419-668-5423  
 Web address: www.huroncohealth.com

**RECOMMENDED ACETAMINOPHEN DOSAGE**

(Tylenol™, Tempra™, non-aspirin)

Child's Age/Weight		Suspension Drops 80 mg/0.8 ml Dropper full	Children's Meltaways 80 mg tabs	Children's Tylenol Suspension Liquid 160 mg/5 ml	Junior Tylenol Meltaways 160 mg
0-3 mos.	6-11 lbs	½ dropper (0.4 ml)*			
4-11 mos.	12-17 lbs.	1 dropper (0.8 ml)**			
12-23 mos.	18-23 lbs.	1 ½ dropper (1.2 ml)***			
2-3 yrs	24-35 lbs.		2 tablets	1 teaspoon	
4-5 yrs.	36-47 lbs.		3 tablets	1 ½ teaspoon	
5-8 yrs.	48-59 lbs.		4 tablets	2 teaspoons	2 tablets
9-10 yrs.	60-71 lbs.		5 tablets	2 ½ teaspoons	2 ½ tablets
11 yrs.	72-95 lbs.		6 tablets	3 teaspoons	3 tablets
12-14 yrs.	96 lbs.				4 tablets

\* 0.4 ml = smallest line on dropper (Use only the dropper or cup that come with that medication bottle)

\*\* 0.8 ml = highest line on dropper

\*\*\* 1.2 ml = highest line on dropper plus smallest line on dropper

**RECOMMENDED IBUPROFEN DOSAGE**

(Motrin™)

Child's Age/Weight		Oral Drops 50 mg/1.25 ml	Suspension 100 mg/5 ml	Chewable Tablets 50 mg	Caplets Tablets 100 mg	Caplets 100 mg
6-11 mos.	12-17 lbs.	½ to 1 dropper	¼ to ½ teaspoon			
12-23 mos.	18-23 lbs.	1 to 2 dropper	½ to 1 teaspoon	1 to 2 tabs	½ to 1 tab	
2-3 yrs	24-35 lbs.	2 droppers	1 teaspoon	1 ½ to 3 tabs	¾ to 1 ½ tab	
4-5 yrs	36-47 lbs.		1 ½ teaspoons	2 to 4 tabs	1 to 2 tabs	
6-8 yrs.	48-59 lbs.		2 teaspoons	2 ½ to 5 tabs	1 ¼ to 2 ½ tabs	2 caplets
9-10 yrs.	60-71 lbs.		2 ½ teaspoons	3 to 6 tabs	1 ½ to 3 tabs	2 ½ caps
11 yrs	72-95 lbs.		3 teaspoons	4 to 8 tabs	2 to 4 tabs	3 caplets

**RECOMMENDED DOSAGE FOR ADVIL**

Weight	Infant Advil 1.25ml 50mg	Age	Children Dosage	Chewable Tablets 50mg	Chewable 100 mg
12-17lbs.	1.25 ml	6-11 mths			
18-23 lbs.	1.875 ml	12-23 mths			
Under 24 lbs.		Under 2 yrs.	Ask a Doctor	Ask a Doctor	
24 to 35 lbs.		2 to 3 yrs.	1 teaspoon	2 tablets	
36 to 47 lbs.		4 to 5 yrs.	1 ½ teaspoon	3 tablets	
48 to 59 lbs.		6 to 8 yrs.	2 teaspoons	4 tablets	2 tablets
60 to 71 lbs.		9 to 10 yrs.	2 ½ teaspoons	5 tablets	2 ½ tablets
72 to 95 lbs.		11 years	3 teaspoons	6 tablets	3 tablets

**\*\*NOTE:** One dose lasts 6 to 8 hours\*\*

**DO NOT GIVE ASPIRIN! These medications contain Aspirin**

(A Partial List)


Nonprescription Products		Prescription Products		
Alka Seltzer (all flavors)	Anacin	Acuprin 81	Aggrenox	Butalbital
Ascriptin	Bayer Aspirin Products	Carisoprodol	Darvon	Disalcid
Bicarbonate Powders(read labels)		Easprin	Endodan	Equagesic
Bufferin	Doan's Pills	Fiorinal	Fiorital	Gelpirin
Dristan	Ecotrin	Halfprin	Lobac	Lortab
Excedrin	Goody's Headache Powders	Magan	Magsal	Methocarbamol
Norwich Aspirin	Pamprin	Mono-Gesic	Norgesic	Percodan
Pepto-Bismol	Sine-Off	Propoxyphene	Robaxisal	Roxiprin
St. Joseph Adult Aspirin		Salflex	Salsalate	Soma
Vanquish	Ysp Aspirin	Synalgos-DC	Talwin	Trilisate

\*If you are not sure about the medication containing aspirin ask the pharmacist. \*



## After the Shots...

### Frequently Asked Questions?



Your child may need extra love and care after getting immunized. Many of the shots that protect children from serious diseases can also cause discomfort for a while. If you don't find the answers to your questions, call the clinic!

Clinic phone #: **419-668-1652 Ext. 252** or **1-888-694-2443 Ext. 252**

\*Baby & Child immunizations are available by appointment\*

**My child has been fussy since you immunized him/her. What should I do?** After immunization, children may be fussy due to pain and/or fever. You may want to give your child acetaminophen, a medicine that helps to reduce pain and fever. Some examples of acetaminophen are: **Tylenol, Panadol, and Tempra.**

**DO NOT GIVE ASPIRIN.** See chart on back. If the fussiness lasts for more than 24 hours, you should call your doctor / ER.

**My child's arm or leg is swollen, hot and red. What should I do?**

- A clean, cool washcloth may be applied over the sore area.
- If there is more redness or tenderness after 24 hours, call your doctor / ER.
- For pain, give acetaminophen. See chart on back. **DO NOT GIVE ASPIRIN.**

**I think my child has a fever. What should I do?**

Check your child's temperature to find out if there is a fever. Do this by taking an ear or underarm temp. If your child's fever is 102°F or higher, you need to call your doctor/ER.

**Here are some things you can do to reduce a fever:**

- Give your child plenty to drink,
- Clothe your child lightly. **DO NOT** cover or wrap your child tightly!
- Give your child acetaminophen. **DO NOT GIVE ASPIRIN.**
- Sponge your child in a few inches of lukewarm (not cold) bath water.

**My child seems really sick. Should I call the doctor?**

If you are worried **AT ALL**, about how your child looks or feels, please call your doctor/ER.

Call your doctor/ER if you answer "YES" to any of the following questions:	
<input checked="" type="checkbox"/> If yes!	
	Does your child have an ear or underarm temperature of 102 degrees or higher? (You should call the clinic if you are concerned about these temperatures.)
	Is your child pale or limp?
	Has your child been crying for over 3 hours and just won't quit?
	Does your child have a strange cry that isn't normal (a high-pitched cry)?
	Is your child's body shaking, twitching or jerking?
<b>**Notify your doctor/ER/Health Department if any of the above happens to your child. **</b>	